General Disclaimer

One or more of the Following Statements may affect this Document

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.

- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.

- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.

- This document is paginated as submitted by the original source.

- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

Produced by the NASA Center for Aerospace Information (CASI)

"Made available under NASA sponsorship in the interest of early and wide dissemination of Earth Resources Survey Program information and without liability for any use made thereof."

E7.7-10.0.25
CR-149128

Title of Investigation: 28990

Investigation of Environmental Change Pattern in JAPAN

Principal Investigator: Takakazu MARUYASU
Science University of Tokyo
Noda City, Chiba ken, 278, JAPAN

October 29, 1976

Quarterly Progerss Report for period

July-Sept. 1976

(E77-10025) INVESTIGATION OF ENVIRONMENTAL CHANGE PATTERN IN JAPAN Quarterly Progress Report, Jul. - Sep. 1976 (Science Univ. of Tokyo (Japan).) 18 p HC A02/MF A01 CSCL 04B

N77-11488

Unclas
G3/43 00025

Name and Address of National Sponcering Agency: Science and Technology Agency
Kasumigaseki 2-2-1 Chiyodaku
Tokyo, 100, JAPAN

NOV 1976
RECEIVED
NASA STI FACILITY
INPUT BRANCH

RECEIVED BY
NASA STI FACILITY
DATE:
DCAF NO. 721002
PROCESSED BY
☑ NASA STI FACILITY
☐ ESA - SDS ☐ AIAA

Original photography may be purchased from:
EROS Data Center
10th and Dakota Avenue
Sioux Falls, SD 57198

Principal Investigator: Takakazu MARUYASU
Science University of Tokyo

Co-Investigator: Hiroaki OCHIAI
Toba Merchant Marine College

Yasuhiro SUGIMORI
Chief of Disaster Analysis Laboratory,
National Research Center for Disaster Protection
Science and Technology Agency

Daitaro SHOJI
Chief ofHydrographer,Hydrographic Department
Marine Safety Agency

Kaname TAKEDA
Leader of Research Group,National Institute
of Resources.

Kiyoshi TSUCHIYA
National Space Development Agency of JAPAN

Iwao NAKAJIMA
Section Chief of Management Technology,Forest
Experiment Station,Forest Agency

Takamasa NAKANO
Faculty of Science,Tokyo Metropolitan University

Shigechika HAYASHI
Chief of Soil Improvement Laboratory,Hokkaido
National Agricultural Experiment Station
Upland Farming Division,Forest Agency

Seiji HORIKAWA
Faculty of Engineering,University of Tokyo

Shunji MURAI
Insitute of Industrial Science,University of
Tokyo

Kantaro WATANABE
Faculty of Marine Science and Technology,
Tokai University

Taisuke WATANABE
Chief of No 2 Resources Section,Tokai Region
Fisheries Research Laboratory,Fishery Agency

ORIGINAL PAGE IS
OF POOR QUALITY

Title of Investigation : 28990

Investigation of Environmental Change Pattern in Japan

Principal Investigator : Takakazu Maruyasu

October 18, 1976

ORIGINAL PAGE IS
OF POOR QUALITY

SUMMARY

SUBJECT: Investigation of Environmental Change Pattern in Japan

OBJECTIVES: The applications and feasibilities of space sensing of the environment and its change in Japan will be investigated with particular focus on the following subjects;

1. Establishment of ecological indicators of environmental disruption through observation of regional vegetation covers and growing conditions.
2. Application of space-acquired data for understanding environmental changes in the costal and offshore zone.

Summary of the results

The study is still going on and the results as intended at the beginning has not been obtained due to partly lack of data and partly due to delay of data. The results so far obtained are summarized below.

Land use

A detailed land use classification for a large urban area of Tokyo was made using MSS digital data. It was found that residental, commercial, industrial, wooded area and grass land can be successfully classified.

Based on the basic researches on land use and vegetation cover classification, an experiment of operational application of Landsat data in an environmental survey was made for a large area of Kanto and Central JAPAN area. In this experiment air-borne data of training area were also added. The result indicates that a regional planning map can be made very effectively. The systems for this operational application is also developed.

Coastal zone and offshore zone

A mesoscale vortex associated with Large Ocean Current Kuroshio which is a rare phenpmenon was firstly recognized visually through the analysis of MSS data. It was found this vortex affects enormously the effluent patterns of river. A majority of investigators agree that this is a mesoscale oceanographical vortex, however there are few investigators taking this vortex as an atmospheric phenomenon.

Based on the Watanabe's theory and technique the same author made every possible efforts to distinguish between oceanographical and atmospheric phenomena.

The areal influence and pattern of " Red Tide " popularly called due to its color in Ise Bay located in Central Japan is found.Through the analysis of four bands data of MSS it is found Band-4 can show partly red tide,however there remains a problem to distinguish it from other water pollution and smog.

An experiment was made to enhance and distinguish sea surface status through photographic processes.

Geophysics and related field

In an attempt to find the applicability of MSS data in the field of volcanology and geology, MSS data covering Active Volcano Sakurajima were analyzed. It was found that lava flowed out at the time of 3 major erruptions which took place in 1779, 1914 and 1946 can be clearly classified.

ORIGINAL PAGE IS
OF POOR QUALITY

The observed values of MSS include the effects of atmospheric scattering and absorption. An attempt is made to numerically evaluate its effects through numerical integration of radiative transfer equation. It is found that the radiance due to atmospheric effect popularly called as path radiance for Bands 4,5,6 and 7 of MSS amounts as large as 47, 34, 24 and 20% of the measured value (total radiance) respectively.



Fig.1. Theme Extraction Imagery of Kanto Area

Blue : Forest
Yellow : Firm field
White : Bare soil
Red : Urban area, Polluted water




Fig.2. Color Composit Imagery of Westside Tokyo by LANDSAT-1 CCT, used for the determination of the classification and the training area.




Fig.3. Digital Pattern Recognition Output of Fig.2 area.
Classification to 10 items.




Fig.4. Color Composit Imagery of Middle Part of Japan, at the end of September the rice field harvested already.




Fig.5. Land Use Map of Fig.4 area.
Green : Rice field
Yellow : Other crops
Black : Urban
Red : Under construction



Fig.6. Pattern Classification in Mountainous Forest.
Green : Ever green grass land
Brown : Artificial conifer
Yellow : Pine and other conifer

ORIGINAL PAGE IS
OF POOR QUALITY




Fig.7. Land Use Pattern Recognition of North Suburb Tokyo.
Green : Forest
Yellow : Firm field
Red : Urban
Blue : Water




Fig.17. Forest Type Quadrate Map of North Kanto.
5 classes by the tree cover densities. (2x2 km unit)
Total area : 7,460 km^2.




Fig.18. Timber Productivity Classification.
Produced by LANDSAT data and census data.




Fig.19. Recreation Area Zoning in 3 classes.
Calculated from the population data and green capacity of each quadrate unit, also relative distance.




Fig.20. Forest Conservation Planning Map.
Each quadrate unit was evaluated by the potential power for watershade, erosion control, flood control and timber productivity.

ORIGINAL PAGE IS
OF POOR QUALITY

Land Use around Central Tokyo

These 4 photos show the results of land use analysis of Central Tokyo by using the LANDSAT CCT on 14-DEC-72.

A



C



B



D



A: Central Tokyo

Red:Densely built up area
Pink:Residential area
Green:Park and woody field and forest
Light blue:Fresh water
Dark blue: Salty water
Black:Uncategorized area

B: Urban of Tokyo

Red:Built up area
Pink:Residential area

C: Urban of Tokyo

Green:Woody forest as imperial palace
Yellowish green:Grass field as golf links

D: Tokyo bay

Water area divided into sea area and fresh water

ORIGINAL PAGE IS OF POOR QUALITY

Table 1. Land use ratio in the Central Tokyo

NO. OF ORIGINAL VARIABLES = 4
REJECTION LEVEL = 3.00

GRP NO.	CATEGORY	NOB	PERCENT OF TOTAL	ACRES	SQ. KM.
0	UNCATEGORIZED	1570	8.25	1755.10	7.10
1	KOKYO GREEN	1366	7.17	1527.04	6.18
2	SHINBASHI	7799	40.96	8718.46	35.28
3	TODOROKI PINK	6636	34.85	7418.35	30.02
4	ARAKAWA LB	805	4.23	899.91	3.64
5	TOKYO BAY	864	4.54	965.86	3.91
TOTALS		19040	100.00	21284.73	86.14

Table 2. Training area selected for MCP analysis

18-MAR-76 18:15:21

DN	GN	COLOR	NOB	X1	Y1	X2	Y2	X3	Y3	X4	Y4	NAME
0	0	000										
1	1	050	95	1079	80	1094	81	1090	91	1081	90	KOKYO GREEN
2	1	050	9	1051	98	1062	100	1059	101	1055	103	JINGU GREEN
3	1	050	31	1009	105	1018	104	1018	114	1009	111	MEIJI JINGU G
4	2	500	18	1093	129	1103	129	1103	132	1093	132	SHINBASHI
5	2	500	120	1118	67	1131	67	1131	78	1118	78	NIHONBASHI
7	3	544	24	978	196	986	195	985	206	978	205	TODOROKI PINK
8	3	544	39	938	135	951	127	950	133	944	139	CHOOFU PINK
9	3	544	109	872	58	881	59	886	70	870	69	MITAKA PINK
10	4	007	21	1238	73	1242	73	1242	80	1237	80	ARAKAWA LB
11	4	007	159	1253	136	1265	139	1264	158	1257	158	YUMENOSHIMA L B
12	5	003	250	1240	198	1252	201	1251	220	1238	226	TOKYO BAY
13	5	003	333	1164	191	1192	195	1191	214	1167	202	TOKYO HARBOR D B
15	1	050	38	822	360	842	364	836	365	826	367	TAMAHILL-2
16	1	050	30	913	237	925	242	918	244	912	243	TAMA-HILL 3
20	1	050	35	824	361	832	359	835	365	825	366	YOKOHAMA GREEN

Table 3. Categorization table (showing the precision of analysis)

TNG SET	PERCENT CATEGORIZED AS GROUP 0	1	2	3	4	5
1	2.	93.	0.	5.	0.	0.
2	0.	100.	0.	0.	0.	0.
3	3.	97.	0.	0.	0.	0.
4	0.	0.	100.	0.	0.	0.
5	1.	0.	94.	3.	3.	0.
7	0.	0.	0.	100.	0.	0.
8	3.	3.	0.	95.	0.	0.
9	4.	4.	3.	90.	0	0.
10	0.	0.	0.	0.	100.	0.
11	2.	0.	2.	0.	96.	0.
12	2.	0.	0.	0.	0.	98.
13	3.	0.	0.	0.	0.	97.
15	3.	97.	0.	0.	0.	0.
16	0.	97.	0.	3.	0.	0.
20	3.	94.	0.	3.	0.	0.

ORIGINAL PAGE IS OF POOR QUALITY

Land Use




Table 1. LANDSAT data application diagram for Forest conservation plan.

ORIGINAL PAGE IS
OF POOR QUALITY

A Vortex off Shionomisaki (1)




Fig.3

Photograph






Fig.4




Fig.1




Fig.5




Fig.2




Fig.6

Figure's captions are found in page 10.

ORIGINAL PAGE IS
OF POOR QUALITY

A Vortex off Shionomisaki (2)




Fig.7




Fig.9




Fig.10




Fig.8

ORIGINAL PAGE IS
OF POOR QUALITY

Figure's captions of "A vortex off Shionomisaki", coastal zone and offshore zone.

Photograph : Elliptical Vortex off Shiono Misaki, Sep. 11 1975

Fig.1 : Surface Current around the Place where vortex is found by GEK, Aug. 14 - Sep. 3, 1975

Fig.2 : Surface Current by GEK, Sep. 8 - 16, 1975.

Fig.3 : Hypothetical Stream Line of Kuroshio

Fig.4 : Transparency Distribution (m), Sep. 11 and 12, 1975

Fig.5 : Water Color Distribution, Sep. 11 and 12, 1975

Fig.6 : Surface Salinity (%), Sep. 11 and 12, 1975

Fig.7 : Kuroshio and Associated Coastal Current, Sep. 1975

Fig.8 : The Relation Ship between the values of Band 4 and that of 5

Fig.9 : Computer Output of Band 4 near mouth of Tenryu River

Fig.10 : Computer Output of Band 4 near mouth of Kumano River

ORIGINAL PAGE IS
OF POOR QUALITY



Fig.1

Figure's captions are found in page 13.




FIG. 2

Fig.2




FIG. 3

Fig.3

ORIGINAL PAGE IS OF POOR QUALITY




Fig.4




Fig.5




Fig.6




Fig.7

ORIGINAL PAGE IS
OF POOR QUALITY




Fig.8

Figure's Captions

Fig.1 : Landsat-2 MSS Image Covering Ise Bay and Nagoya District.

Fig.2 : Reclaimed areas at Northern part of Yokkaich and Southern part of Nagoya ports which are located to the western part of Ise Bay.

Fig.3 : Reclaimed area in the western part of Nagoya port


Fig.4-7: Computer output of MSS-4, 5, 6 and 7 showing the coastal areas of Ise Bay

Fig.8 : Computer output of MSS-4, 5, 6 and 7 showing Nagoya port

A



B



Sakurajima Island image is extracted from LANDSAT-1 CCT on 2-DEC-72. The area of the naked lava is divided from another area, where the vegetation has been grown up already. Furthermore two types of the lava are classified. One flowed out in 1914 and another by the big erruption in 1946. There are some area where the two classified are mixed.

A: Green; Vegetation grown up area
Brown; Naked lava

B: White; Showa lava (year 1946)
Light brown; Taisho lava (year 1914)

ORIGINAL PAGE IS
OF POOR QUALITY




Fig. Geo-1 Schematic representation of the model. used for the computation.

	$\lambda = 0.55\mu$		$\lambda = 0.65\mu$	
3 Layers	I_D/I_T	I_P/I_T	I_D/I_T	I_P/I_T
	0.534	0.466	0.659	0.341
	$\lambda = 0.75\mu$		$\lambda = 0.95\mu$	
3 Layers	0.762	0.238	0.805	0.195

Table Geo-1 I_D: Intrinsic radiance of the surface

I_P: Path Radiance I_T: Total Radiance

Each value is obtained through numerical integration of Radiative Transfer Equation.

ORIGINAL PAGE IS OF POOR QUALITY